Operator product expansion analysis of Quasi-PDF

BY LUCHANG JIN UConn/RBRC

PKU ITPseminar

10 May 2018

PKU Physics South 408

Talk is mostly based on work "Factorization Theorem Relating Euclidean and Light-Cone Parton Distributions" by Taku Izubuchi, Xiangdong Ji, Luchang Jin, Iain Stewart, and Yong Zhao. arXiv:1801.03917.

- PDF
- Quasi-PDF
- Matching between Quasi-PDF and PDF
- Matching in coordinate space
- Moments and Quasi-PDF
- Renormalization of quasi-PDF
- Conclusion



Deep inelastic scattering (DIS)

large $Q^2 = -g_{\mu,\nu}q^{\mu}q^{\nu}$ limit.

The DIS process probes the structure of the hadron. In particular, the total cross section of DIS can be factorized into the product of PDF and a perturbative calculable hard part.

A intuitive interpretation of PDF is that it describes the **probability** distribution of the **momentum fraction** carried by a particular kind of parton within a **fast moving** hadron.

The parton can be either a quark or a gluon.

PDF is a necessary input to almost all theory predictions for hadron colliders. Lattice QCD is the only way to compute PDF from the first principles. Conventionally, PDF is obtained by fitting some of the experimental data.

Define PDF

$$O(\xi^{-}) = \bar{\psi}(\xi^{-})\gamma^{+}U(\xi^{-},0)\psi(0)$$
(1)

where $U(\xi^-, 0)$ is the gauge link along the light cone direction, $\xi^- = (t-z)/\sqrt{2}$ is the light cone coordinate and t is physical time, and $\gamma^+ = (\gamma^0 + \gamma^z)/\sqrt{2}$.

$$Q\left(\zeta = P^{+}\xi^{-}, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \frac{1}{2P^{+}} \langle P|O(\xi^{-})|P\rangle$$
(2)

 $|P\rangle$ represent a single hadron state with its momentum along z direction. We will call $\zeta = P^+\xi^-$ the loffe time. Renormalization is needed to evaluate the matrix elements. One can conveniently choose $\overline{\text{MS}}$ renormalization scheme and denote the renormalization scale to be μ . Another scale Λ_{QCD} is introduced.

$$q\left(x,\frac{\mu}{\Lambda_{\rm QCD}}\right) = \int \frac{d\zeta}{2\pi} e^{-ix\zeta} Q\left(\zeta,\frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(3)

Finally, PDF of the hadron is defined above as $q(x, \mu / \Lambda)$. This definition has a special property:

$$\bar{q}(x,\mu/\Lambda_{\rm QCD}) = -q(-x,\mu/\Lambda_{\rm QCD})$$
 (4)

4/32

Define PDF

$$F_1(x, Q^2) = \int \frac{d\xi}{|\xi|} C_1\left(\frac{x}{\xi}, \frac{Q^2}{\mu^2}\right) q\left(\xi, \frac{\mu}{\Lambda_{\rm QCD}}\right) + \text{higher twist contributions}$$
(5)

where $F_1(x, Q^2)$ is commonly known as the deep-inelastic scattering (DIS) structure function. Operator product expansion (OPE)

$$O(\xi^{-}) = \sum_{n} \frac{(i\xi^{-})^{n}}{n!} O^{+\dots+}$$
(6)

where $O^{+...+}$ is a component of the following symmetric traceless operator:

$$O^{\mu_0 \mu_1 \dots \mu_n} = \bar{\psi} \gamma^{(\mu_0} i D^{\mu_1} \dots i D^{\mu_n)} \psi - \text{trace}$$
(7)

The moments of PDF $a_n(\mu/\Lambda)$ is related with OPE:

$$\frac{1}{2} \langle P | O^{\mu_0 \mu_1 \dots \mu_n} | P \rangle = a_{n+1} \left(\frac{\mu}{\Lambda_{\text{QCD}}} \right) \left(P^{\mu_0} P^{\mu_1} \dots P^{\mu_n} - \text{trace} \right)$$
(8)

Moments of PDF

Here we demonstrate the reason we call $a_n(\mu/\Lambda_{\rm QCD})$ the moment of PDF.

On one hand:

$$\frac{1}{2} \langle P | O^{+\dots+} | P \rangle = a_{n+1} \left(\frac{\mu}{\Lambda_{\text{QCD}}} \right) (P^+)^{n+1}$$
(9)

$$Q\left(\zeta = P^{+}\xi^{-}, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \sum_{n} \frac{(i\zeta)^{n}}{n!} a_{n+1}\left(\frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(10)

$$\left(\frac{\partial}{\partial i \zeta}\right)^{n} Q\left(\zeta, \frac{\mu}{\Lambda_{\rm QCD}}\right)\Big|_{\zeta=0} = a_{n+1}\left(\frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(11)

On the other hand:

$$Q\left(\zeta, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \int dx \, e^{ix\zeta} q\left(x, \frac{\mu}{\Lambda_{\rm QCD}}\right) \tag{12}$$

$$\left(\frac{\partial}{\partial i \zeta}\right)^{n} Q\left(\zeta, \frac{\mu}{\Lambda_{\rm QCD}}\right)\Big|_{\zeta=0} = \int dx \, x^{n} \, q\left(x, \frac{\mu}{\Lambda_{\rm QCD}}\right) \tag{13}$$

Talk is mostly based on work "Factorization Theorem Relating Euclidean and Light-Cone Parton Distributions" by Taku Izubuchi, Xiangdong Ji, Luchang Jin, Iain Stewart, and Yong Zhao. arXiv:1801.03917.

- PDF
- Quasi-PDF
- Matching between Quasi-PDF and PDF
- Matching in coordinate space
- Moments and Quasi-PDF
- Renormalization of quasi-PDF
- Conclusion

Calculate PDF on lattice

$$O^{\mu_0\mu_1...\mu_n} = \bar{\psi}\gamma^{(\mu_0}i D^{\mu_1}...i D^{\mu_n)}\psi - \text{trace}$$

Moments of PDF

In principle, the moments of PDF can be calculated on lattice.

There are extensive lattice studies of moments of PDF.

However, because the lattice regulator breaks rotational symmetry, the operator $O^{\mu_0\mu_1...\mu_n}$ $(n \ge 4)$ mixes with lower dimensional operator. It is very hard to compute the 4th or higher order moments directly.

Calculate PDF on lattice

$$O^{\mu_0\mu_1...\mu_n} = \bar{\psi} \gamma^{(\mu_0} i D^{\mu_1}... i D^{\mu_n)} \psi - \text{trace}$$

Non-local operator

Using a non-local operator (or two local operators) does not have the mixing problem. The new problem is: in Euclidean space, non-local implies non-zero z^2 , here z is the 4-vector characterizing the shape of the non-local operator, but the PDF is related to the light-cone region where $z^2 = 0$. Non-zero z^2 usually implies higher twist effects. The issue then becomes how to suppress these higher twist effects.

Quasi-PDF: large momentum hadron state

"Parton Physics on a Euclidean Lattice" by Xiangdong Ji, Phys.Rev.Lett. 110 (2013) 262002

Other approaches

- "Deep-inelastic scattering and the operator product expansion in lattice QCD" by William **Detmold** and **David Lin**, 2006.
- "Restoration of rotational symmetry in the continuum limit of lattice field theories" by Zohreh Davoudi and Martin Savage, Phys.Rev. D86 (2012) 054505.
- "Parton Distribution Function from the Hadronic Tensor on the Lattice" by Keh-Fei Liu, 2016.
- "Nucleon structure functions from lattice operator product expansion" by **QCDSF**, 2017.

Define Quasi-PDF

Concept is introduced by **Xiangdong Ji** Phys.Rev.Lett. 110 (2013) 262002. Introduce a non-local operator which is:

- Calculable on Euclidean spacetime lattice.
- Similar to the operator used to define PDF.

$$\tilde{O}(z) = \bar{\psi}(z)\gamma^z U(z,0)\psi(0) \tag{14}$$

Then we define its matrix elements and Fourier transformation similarly.

$$\tilde{Q}\left(\zeta = P^{z} z, \mu^{2} z^{2} = \frac{\mu^{2} \zeta^{2}}{(P^{z})^{2}}, \frac{\mu}{\Lambda_{\text{QCD}}}\right) = \frac{1}{2 P^{z}} \langle P | \tilde{O}(z) | P \rangle$$
(15)

Note an important difference compare with PDF is the introduction of z^2 .

$$\tilde{q}\left(\tilde{x}, \frac{\mu^2}{(P^z)^2}, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \int \frac{d\zeta}{2\pi} e^{-i\tilde{x}\zeta} \tilde{Q}\left(\zeta, \frac{\mu^2 \zeta^2}{(P^z)^2}, \frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(16)

The hope is: the quasi-PDF $\tilde{q}(\tilde{x}, \mu^2/(P^z)^2, \mu/\Lambda_{QCD})$ can be used to determine PDF if the momentum of the hadron P^z is large.

PDF notations

11/32

Summarize the notations so far:

PDF (light-cone)Quasi-PDFMomemtum fraction x - space $q\left(x, \frac{\mu}{\Lambda_{QCD}}\right)$ $\tilde{q}\left(x, \frac{\mu^2}{(P^z)^2}, \frac{\mu}{\Lambda_{QCD}}\right)$

 $\text{loffe-time } \zeta \text{ - space} \qquad Q\Big(\zeta = P^+ \xi^-, \frac{\mu}{\Lambda_{\text{QCD}}}\Big) \quad \tilde{Q}\Big(\zeta = P^z z, \, \mu^2 z^2 = \frac{\mu^2 \zeta^2}{(P^z)^2}, \frac{\mu}{\Lambda_{\text{QCD}}}\Big)$

Coordinate space and momentum fraction \boldsymbol{x}

$$q\left(x,\frac{\mu}{\Lambda_{\rm QCD}}\right) = \int \frac{d\zeta}{2\pi} e^{-ix\zeta} Q\left(\zeta,\frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(17)

$$\tilde{q}\left(\tilde{x}, \frac{\mu^2}{(P^z)^2}, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \int \frac{d\zeta}{2\pi} e^{-i\tilde{x}\zeta} \tilde{Q}\left(\zeta, \frac{\mu^2\zeta^2}{(P^z)^2}, \frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(18)

Different Fourier transformation:

$$\mathcal{P}\left(x,\mu^2 z^2,\frac{\mu}{\Lambda_{\rm QCD}}\right) = \int \frac{d\zeta}{2\pi} e^{-ix\zeta} \tilde{Q}\left(\zeta,\mu^2 z^2,\frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(19)

Talk is mostly based on work "Factorization Theorem Relating Euclidean and Light-Cone Parton Distributions" by Taku Izubuchi, Xiangdong Ji, Luchang Jin, Iain Stewart, and Yong Zhao. arXiv:1801.03917.

- PDF
- Quasi-PDF
- Matching between Quasi-PDF and PDF
- Matching in coordinate space
- Moments and Quasi-PDF
- Renormalization of quasi-PDF
- Conclusion

Matching between Quasi-PDF and PDF

 $\tilde{O}(z) = \bar{\psi}(z)\gamma^z U(z,0)\psi(0)$

OPE can be performed for quasi-PDF operator ($z^2 \rightarrow 0$):

$$\tilde{O}(z) = \sum_{n} C_n(\mu^2 z^2) \frac{(i z)^n}{n!} O^{z \cdots z} + \text{higher twist contributions}$$
(20)

Similar to $O^{+\cdots+}$, $O^{z\cdots z}$ is also a component of the operator:

$$O^{\mu_0\mu_1\dots\mu_n} = \bar{\psi}\gamma^{(\mu_0}iD^{\mu_1}\dots iD^{\mu_n)}\psi - \text{trace}$$

$$\frac{1}{2}\langle P|O^{z\dots z}|P\rangle = a_{n+1}\left(\frac{\mu}{\Lambda_{\text{QCD}}}\right)((P^z)^{n+1} - \text{trace})$$
(21)

13/32

Therefore

$$\tilde{Q}\left(\zeta = P^{z}z, \mu^{2}z^{2}, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \sum_{n} C_{n}(\mu^{2}z^{2}) \frac{(i\zeta)^{n}}{n!} a_{n+1}\left(\frac{\mu}{\Lambda_{\rm QCD}}\right) + \text{higher twist}$$
(22)

Comparing OPE of Quasi-PDF and PDF

$$O(\xi^{-}) = \sum_{n} \frac{(i\xi^{-})^{n}}{n!} O^{+\dots+}$$
(23)

14/32

$$\tilde{O}(z) = \sum_{n} C_n(\mu^2 z^2) \frac{(i z)^n}{n!} O^{z \cdots z} + \text{higher twist contributions}$$
(24)

 $O^{+\cdots+}$, $O^{z\cdots z}$ are components of the symmetric traceless operator:

$$O^{\mu_0\mu_1...\mu_n} = \bar{\psi} \gamma^{(\mu_0} i D^{\mu_1}... i D^{\mu_n)} \psi - \text{trace}$$
(25)

The matrix elements are:

$$\frac{1}{2} \langle P | O^{+\dots+} | P \rangle = a_{n+1} \left(\frac{\mu}{\Lambda_{\text{QCD}}} \right) (P^+)^{n+1}$$
(26)

$$\frac{1}{2} \langle P | O^{z \dots z} | P \rangle = a_{n+1} \left(\frac{\mu}{\Lambda_{\text{QCD}}} \right) \left((P^z)^{n+1} - \text{trace} \right)$$
(27)

Use large momentum P^z to suppress the higher twist and trace contributions.

$$\tilde{Q}\left(\zeta = P^{z}z, \frac{\mu^{2}\zeta^{2}}{(P^{z})^{2}}, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \sum_{n} C_{n}\left(\frac{\mu^{2}\zeta^{2}}{(P^{z})^{2}}\right) \frac{(i\,\zeta)^{n}}{n!} a_{n+1}\left(\frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(28)

Matching between Quasi-PDF and PDF

$$\tilde{Q}\left(\zeta, \frac{\mu^2 \zeta^2}{(P^z)^2}, \frac{\mu}{\Lambda_{\text{QCD}}}\right) = \sum_n C_n\left(\frac{\mu^2 \zeta^2}{(P^z)^2}\right) \frac{(i\zeta)^n}{n!} a_{n+1}\left(\frac{\mu}{\Lambda_{\text{QCD}}}\right)$$
$$= \sum_n C_n\left(\frac{\mu^2 \zeta^2}{(P^z)^2}\right) \frac{(i\zeta)^n}{n!} \int_{-1}^1 dx \, x^n \, q\left(x, \frac{\mu}{\Lambda_{\text{QCD}}}\right)$$

15/32

$$\begin{split} \tilde{q}\bigg(\tilde{x}, \frac{\mu^2}{(P^z)^2}, \frac{\mu}{\Lambda_{\rm QCD}}\bigg) &= \int \frac{d\,\zeta}{2\pi} e^{-i\tilde{x}\zeta} \tilde{Q}\bigg(\zeta, \frac{\mu^2 \zeta^2}{(P^z)^2}, \frac{\mu}{\Lambda_{\rm QCD}}\bigg) \\ &= \int_{-1}^1 d\,x \bigg[\int \frac{d\,\zeta}{2\pi} e^{-i\tilde{x}\zeta} \sum_n C_n\bigg(\frac{\mu^2 \zeta^2}{(P^z)^2}\bigg) \frac{(i\,\zeta)^n}{n!} x^n \bigg] q\bigg(x, \frac{\mu}{\Lambda_{\rm QCD}}\bigg) \end{split}$$

Define:

$$C\left(\frac{\tilde{x}}{x},\frac{\mu^2}{(xP^z)^2}\right) = \int \frac{d(x\zeta)}{2\pi} e^{-i\frac{\tilde{x}}{x}(x\zeta)} \sum_n C_n\left(\frac{\mu^2(x\zeta)^2}{(xP^z)^2}\right) \frac{(ix\zeta)^n}{n!}$$
(29)

We have:

$$\tilde{q}\left(\tilde{x},\frac{\mu^2}{(P^z)^2},\frac{\mu}{\Lambda}\right) = \int_{-1}^1 \frac{dx}{|x|} C\left(\frac{\tilde{x}}{x},\frac{\mu^2}{(xP^z)^2}\right) q\left(x,\frac{\mu}{\Lambda}\right)$$
(30)

Determine the matching kernel

The same matching formula should also apply to quark state (quasi-)PDF. We denote the quark state (quasi-PDF) by the same function name but the argument μ/Λ_{QCD} absent.

$$\tilde{q}\left(\tilde{x}, \frac{\mu^2}{(P^z)^2}\right) = \int_{-1}^1 \frac{dx}{|x|} C\left(\frac{\tilde{x}}{x}, \frac{\mu^2}{(x P^z)^2}\right) q(x)$$
(31)

One can perturbatively compute q(x) and $\tilde{q}(\tilde{x}, \mu^2/(P^z)^2)$, then solves the matching kernel C.

Using pure dimensional regularization (canceling UV and IR poles with $\epsilon_{\rm UV} = \epsilon_{\rm IR}$), massless quark, the quark state PDF does not depend on any quantity that has a scale, therefore all the loop contribution will be zero and it is equal to the tree level result:

$$q(x) = \delta(x-1) \tag{32}$$

Immediately we have (in \overline{MS} scheme):

$$C\left(\tilde{x}, \frac{\mu^2}{(P^z)^2}\right) = \tilde{q}\left(\tilde{x}, \frac{\mu^2}{(P^z)^2}\right)$$
(33)

One loop calculation for $C(\tilde{x}, \mu^2/(P^z)^2)$ has already been done.

16/32

Fourier transform differently

Quasi-PDF is defined to be a Fourier transform of the coordinate space matrix elements:

$$\tilde{q}\left(\tilde{x}, \frac{\mu^2}{(P^z)^2}, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \int \frac{d\zeta}{2\pi} e^{-i\tilde{x}\zeta} \tilde{Q}\left(\zeta, \frac{\mu^2 \zeta^2}{(P^z)^2}, \frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(34)

A.V. Radyushkin discovered that one can perform a slightly different Fourier transformation, he call the resulting quantity **Pseudo-PDF** (which is only non-zero when $|x| \leq 1$):

$$\mathcal{P}\left(x,\mu^2 z^2,\frac{\mu}{\Lambda_{\rm QCD}}\right) = \int \frac{d\zeta}{2\pi} e^{-ix\zeta} \tilde{Q}\left(\zeta,\mu^2 z^2,\frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(35)

Similarly, we can prove the matching formula using OPE:

$$\mathcal{P}\left(x,\mu^2 z^2,\frac{\mu}{\Lambda_{\rm QCD}}\right) = \int_{-1}^{1} \frac{dx}{|x|} \mathcal{C}\left(\frac{\tilde{x}}{x},\mu^2 z^2\right) q\left(x,\frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(36)

Again, with dimensional regularization and massless quark, the matching kernel is the same as the quark state pseudo-PDF:

$$\mathcal{C}(x,\mu^2 z^2) = \mathcal{P}(x,\mu^2 z^2) = \int \frac{d\zeta}{2\pi} e^{-ix\zeta} \tilde{Q}(\zeta,\mu^2 z^2)$$
(37)

This can be determined perturbatively and one-loop result has been obtained.

Comparing the matching formulae

DIS structure function:

$$F_1(x, Q^2) = \int_{-1}^1 \frac{d\xi}{|\xi|} C_1\left(\frac{x}{\xi}, \frac{Q^2}{\mu^2}\right) q\left(\xi, \frac{\mu}{\Lambda_{\text{QCD}}}\right) + \text{higher twist contribs}$$

18/32

Quasi-PDF:

$$\tilde{q}\left(\tilde{x}, \frac{\mu^2}{(P^z)^2}, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \int_{-1}^1 \frac{dx}{|x|} C\left(\frac{\tilde{x}}{x}, \frac{\mu^2}{(xP^z)^2}\right) q\left(x, \frac{\mu}{\Lambda_{\rm QCD}}\right) + \text{higher twist contribs}$$

Pseudo-PDF:

$$\mathcal{P}\left(\tilde{x}, \mu^2 z^2, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \int_{-1}^1 \frac{dx}{|x|} \mathcal{C}\left(\frac{\tilde{x}}{x}, \mu^2 z^2\right) q\left(x, \frac{\mu}{\Lambda_{\rm QCD}}\right) + \text{higher twist contribs}$$

• The three scales are Q^2 , $(x P^z)^2$, $1/z^2$. Note that they are all related with single parton.

- One may view quasi-PDF and pseudo-PDF as lattice observables which can be "factorized" into PDF. One may find other relevant and perhaps better lattice observables. Yan-Qing Ma and Jian-Wei Qiu, arXiv:1404.6860, Phys.Rev.Lett. 120, 022003 (2018).
- In particular, F₁(x, Q²) can also be computed on lattice. Many different approaches: QCDSF Phys.Rev.Lett. 118 (2017) no.24, 242001; Keh-Fei Liu arXiv:1603.07352;
 William Detmold and David Lin Phys.Rev. D73 (2006) 014501.

Talk is mostly based on work "Factorization Theorem Relating Euclidean and Light-Cone Parton Distributions" by Taku Izubuchi, Xiangdong Ji, Luchang Jin, Iain Stewart, and Yong Zhao. arXiv:1801.03917.

- PDF
- Quasi-PDF
- Matching between Quasi-PDF and PDF
- Matching in coordinate space
- Moments and Quasi-PDF
- Renormalization of quasi-PDF
- Conclusion

Matching in coordinate space

It would be interesting to see what the matching formula looks like in coordinate space.

$$Q\left(\zeta = P^{+}\xi^{-}, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \sum_{n} \frac{(i\,\zeta)^{n}}{n!} a_{n+1}\left(\frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(38)

$$\begin{split} \tilde{Q}\bigg(\zeta = P^{z} z, \mu^{2} z^{2}, \frac{\mu}{\Lambda_{\rm QCD}}\bigg) &= \sum_{n} C_{n}(\mu^{2} z^{2}) \frac{(i \zeta)^{n}}{n!} a_{n+1}\bigg(\frac{\mu}{\Lambda_{\rm QCD}}\bigg) \\ &= \sum_{n} \bigg[\int_{0}^{1} d \, \alpha \, \mathcal{C}(\alpha, \mu^{2} z^{2}) \, \alpha^{n}\bigg] \frac{(i \zeta)^{n}}{n!} a_{n+1}\bigg(\frac{\mu}{\Lambda_{\rm QCD}}\bigg) \\ &= \int_{0}^{1} d \, \alpha \, \mathcal{C}(\alpha, \mu^{2} z^{2}) \sum_{n} \alpha^{n} \frac{(i \zeta)^{n}}{n!} a_{n+1}\bigg(\frac{\mu}{\Lambda_{\rm QCD}}\bigg) \\ &= \int_{0}^{1} d \, \alpha \, \mathcal{C}(\alpha, \mu^{2} z^{2}) \, Q\bigg(\alpha \, \zeta, \frac{\mu}{\Lambda_{\rm QCD}}\bigg) \end{split}$$

where

$$C_n(\mu^2 z^2) = \int_0^1 d\alpha \, \mathcal{C}(\alpha, \mu^2 z^2) \, \alpha^n \tag{39}$$

20/32

Matching in coordinate space

21/32

$$\tilde{Q}\left(\zeta = P^{z} z, \mu^{2} z^{2}, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \int_{0}^{1} d\alpha \, \mathcal{C}(\alpha, \mu^{2} z^{2}) \, Q\left(\alpha \, \zeta, \frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(40)

The form of the matching formula does not imply z is fixed in the matching. In fact, we can easily change the parameter in the above formula:

$$\tilde{Q}\left(\zeta = P^{z} z, \frac{\mu^{2} \zeta^{2}}{(P^{z})^{2}}, \frac{\mu}{\Lambda_{\text{QCD}}}\right) = \int_{0}^{1} d\alpha \mathcal{C}\left(\alpha, \frac{\mu^{2} \zeta^{2}}{(P^{z})^{2}}\right) Q\left(\alpha \zeta, \frac{\mu}{\Lambda_{\text{QCD}}}\right)$$
(41)

- One can use these coordinate space matching formula to derive the matching formula for quasi-PDF.
- Or, one can apply the coordinate space matching formula directly to the lattice data, which are also measured in the coordinate space. One can then Fourier transform the one-loop matched, light-cone, loffe-time distribution Q to obtain PDF.

Talk is mostly based on work "Factorization Theorem Relating Euclidean and Light-Cone Parton Distributions" by Taku Izubuchi, Xiangdong Ji, Luchang Jin, Iain Stewart, and Yong Zhao. arXiv:1801.03917.

- PDF
- Quasi-PDF
- Matching between Quasi-PDF and PDF
- Matching in coordinate space
- Moments and Quasi-PDF
- Renormalization of quasi-PDF
- Conclusion

Moments and Quasi-PDF

The coefficients of the Taylor expansion of Q gives the moments of PDF.

$$Q\left(\zeta = P^{+}\xi^{-}, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \sum_{n} \frac{(i\zeta)^{n}}{n!} a_{n+1}\left(\frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(42)

However, the moments of quasi-PDF, in general, do not exist. (Note that the moments of pseudo-PDF do exist, but require non-trivial matching coefficients, given by $C_n(\mu^2 z^2)$, which we have computed explicitly at one-loop.) Based on OPE:

$$\tilde{Q}\left(\zeta = P^{z} z, \mu^{2} z^{2}, \frac{\mu}{\Lambda_{\text{QCD}}}\right) = \sum_{n} C_{n}(\mu^{2} z^{2}) \frac{(i \zeta)^{n}}{n!} a_{n+1}\left(\frac{\mu}{\Lambda_{\text{QCD}}}\right)$$
(43)

The reason is that C_n is singular in the $z \rightarrow 0$ limit. Especially, for the first moment:

$$\tilde{Q}\left(\zeta = P^{z} z, \mu^{2} z^{2}, \frac{\mu}{\Lambda_{\text{QCD}}}\right)\Big|_{P^{z} z \to 0} = C_{0}(\mu^{2} z^{2})a_{1}\left(\frac{\mu}{\Lambda_{\text{QCD}}}\right) = C_{0}(\mu^{2} z^{2})$$
(44)

In $\overline{\mathrm{MS}}$ scheme at one-loop order:

$$C_0(\mu^2 z^2) = 1 + \frac{\alpha_s C_F}{2\pi} \left[\frac{3}{2} \log \left(\frac{\mu^2 z^2 e^{2\gamma_E}}{4} \right) + \frac{7}{2} \right].$$
(45)

Moments and Quasi-PDF

In the $z \rightarrow 0$ limit (P^z fixed):

$$\tilde{Q}\left(\zeta = P^{z} z, \mu^{2} z^{2}, \frac{\mu}{\Lambda_{\text{QCD}}}\right)\Big|_{z \to 0} = C_{0}(\mu^{2} z^{2}) = 1 + \frac{\alpha_{s} C_{F}}{2\pi} \left[\frac{3}{2} \log\left(\frac{\mu^{2} z^{2} e^{2\gamma_{E}}}{4}\right) + \frac{7}{2}\right] \quad (46)$$

However, one might have expected a different result:

$$\tilde{Q}\left(\zeta = P^{z} z, \mu^{2} z^{2}, \frac{\mu}{\Lambda_{\text{QCD}}}\right)\Big|_{z \to 0} = \frac{1}{2 P^{z}} \langle P | \bar{\psi}(0) \gamma^{z} \psi(0) | P \rangle = 1?$$

$$(47)$$

The singularity is a result of the \overline{MS} scheme subtraction. Before renormalization:

$$\tilde{Q}\left(\zeta = P^{z} z, \mu^{2} z^{2}, \frac{\mu}{\Lambda_{\text{QCD}}}\right)\Big|_{z \to 0} - 1 \sim \frac{(\mu^{2} z^{2})^{\epsilon}}{\epsilon} \to 0$$
(48)

However, if we expand on ϵ first

$$\frac{(\mu^2 z^2)^{\epsilon}}{\epsilon} = \frac{1}{\epsilon} + \log(\mu^2 z^2)$$
(49)

24/32

The result is logarithmic divergent at 1-loop after \overline{MS} subtraction. (Not true for some other scheme.)

Talk is mostly based on work "Factorization Theorem Relating Euclidean and Light-Cone Parton Distributions" by Taku Izubuchi, Xiangdong Ji, Luchang Jin, Iain Stewart, and Yong Zhao. arXiv:1801.03917.

- PDF
- Quasi-PDF
- Matching between Quasi-PDF and PDF
- Matching in coordinate space
- Moments and Quasi-PDF
- Renormalization of quasi-PDF
- Conclusion

Other operators

Light cone:

$$O(\xi^{-}) = \bar{\psi}(\xi^{-})\gamma^{+}U(\xi^{-},0)\psi(0)$$
(50)

Initial version:

$$\tilde{O}(z) = \bar{\psi}(z)\gamma^z U(z,0)\psi(0)$$
(51)

• Use t direction gamma matrix:

$$\bar{\psi}(z)\gamma^t U(z,0)\psi(0) \tag{52}$$

 Two vector current operators Y.Q. Ma, J.W. Qiu (possibly flavor changing W. Detmold, D. Lin):

$$J_{\mu}(z)J_{\nu}(0) \tag{53}$$

26/32

Renormalization of quasi-PDF

$$\tilde{O}(z) = \bar{\psi}(z)\gamma^z U(z,0)\psi(0)$$

This is not the first time we deal with this kind of operator. In heavy quark effective theory (HQET), we also have this kind of operator.

According to X. Ji, J.H. Zhang, Y. Zhao arXiv:1706.08962 and T. Ishikawa, Y.Q. Ma, J.W. Qiu arXiv:1707.03107, the non-local operator renormalize multiplicatively:

$$\left[\tilde{O}(z)\right]^{\overline{\mathrm{MS}}} = Z(\mu^2 z^2, \mu a) \left[\tilde{O}(z)\right]^{\mathrm{Lat}}$$
(54)

However, there is a subtle caveat discovered by **M. Constantinou** and **H. Panagopoulos** arXiv:1705.11193. The above renormalization formula is only correct for Chiral lattice fermion actions, e.g. domain wall fermion (DWF). For some other popular fermion actions like Wilson fermion, there is a mixing:

$$\left[\tilde{O}(z)\right]^{\overline{\mathrm{MS}}} = Z(\mu^2 z^2, \mu a) \left[\tilde{O}(z)\right]^{\mathrm{Lat}} + Z'(\mu^2 z^2, \mu a) \left[\tilde{O}'(z)\right]^{\mathrm{Lat}}$$
(55)

$$\tilde{O}'(z) = \bar{\psi}(z)U(z,0)\psi(0) \tag{56}$$

The mixing is also absent if one use the γ^t operator: $\bar{\psi}(z)\gamma^t U(z,0)\psi(0)$.

Renormalization of quasi-PDF

28/32

In practice, it is convenient to renormalize the lattice operator first and measure the renormalized matrix elements on the lattice. However, it should be noted that the form of the matching formula is unchanged when applied to the bare lattice results, since the initial OPE formula has the same shape:

$$\left[\tilde{Q}\left(\zeta = P^{z}z, \left(\frac{z}{a}\right)^{2}, \frac{1}{a\Lambda_{\rm QCD}}\right)\right]^{\rm Lat} = \sum_{n} \frac{C_{n}(\mu^{2}z^{2})}{Z(\mu^{2}z^{2}, \mu a)} \frac{(i\zeta)^{n}}{n!} a_{n+1}\left(\frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(57)

To avoid computing the renormalization function Z, which depends on the lattice action, one can compute the ratio of two matrix elements using the same lattice operator. The numerator is simply $\left[\tilde{Q}\left(\zeta = P^z \, z, \left(\frac{z}{a}\right)^2, \frac{1}{a \, \Lambda_{\rm QCD}}\right)\right]^{\rm Lat}$, for the denominator, there are two common choices

- RI/MOM: use the matrix elements of a large Euclidean momentum (off-shell) quark state.
 - I. Stewart and Y. Zhao.
- Ratio: Reduced loffe-time distribution (use γ^t): $\left[\tilde{Q}_{\gamma^t}\left(0, \left(\frac{z}{a}\right)^2, \frac{1}{a\Lambda_{\rm QCD}}\right)\right]^{\rm Lat} = \frac{C_0(\mu^2 z^2)}{Z(\mu^2 z^2, \mu a)}$.

A. V. Radyushkin.

The requirement for the denominator is that the corresponding matrix elements in the continuum renormalization scheme can be calculated perturbatively.

Renormalization of quasi-PDF

Mathing formula for the renormalized coordinate space quasi-PDF using the "ratio" method:

$$\tilde{Q}_{R}\left(\zeta = P^{z}z, \mu^{2}z^{2}, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \frac{\tilde{Q}_{\gamma^{t}}\left(\zeta = P^{z}z, \mu^{2}z^{2}, \frac{\mu}{\Lambda_{\rm QCD}}\right)}{\tilde{Q}_{\gamma^{t}}\left(0, \mu^{2}z^{2}, \frac{\mu}{\Lambda_{\rm QCD}}\right)}$$
(58)

$$= \int_{0}^{1} d\alpha \frac{\mathcal{C}_{\gamma_{t}}(\alpha, \mu^{2} z^{2})}{\int_{0}^{1} d\alpha' \mathcal{C}_{\gamma_{t}}(\alpha', \mu^{2} z^{2}) Q\left(0, \frac{\mu}{\Lambda_{\text{QCD}}}\right)} Q\left(\alpha \zeta, \frac{\mu}{\Lambda_{\text{QCD}}}\right) + \mathcal{O}(z^{2} \Lambda_{\text{QCD}}^{2}) \qquad (59)$$
$$= \int_{0}^{1} d\alpha \mathcal{C}_{R}(\alpha, \mu^{2} z^{2}) Q\left(\alpha \zeta, \frac{\mu}{\Lambda_{\text{QCD}}}\right) + \mathcal{O}(z^{2} \Lambda_{\text{QCD}}^{2}) \qquad (60)$$

where

$$\mathcal{C}_R(\alpha, \mu^2 z^2) = \delta(1-x) + \mathcal{C}_R^{(1)}(x, \mu^2 z^2) + \cdots$$
(61)

$$\mathcal{C}_{R}^{(1)}(x,\mu^{2}z^{2}) = \frac{\alpha_{s}C_{F}}{2\pi}\theta(x)\theta(1-x) \\
\times \left(\frac{1+x^{2}}{1-x}\left[-\log(\mu^{2}z^{2}) - \log\left(\frac{e^{2\gamma_{E}}}{4}\right) - 1\right] - \frac{4\log(1-x)}{1-x} + 2(1-x)\right)_{+}$$
(62)

29/32

Talk is mostly based on work "Factorization Theorem Relating Euclidean and Light-Cone Parton Distributions" by Taku Izubuchi, Xiangdong Ji, Luchang Jin, Iain Stewart, and Yong Zhao. arXiv:1801.03917.

- PDF
- Quasi-PDF
- Matching between Quasi-PDF and PDF
- Matching in coordinate space
- Moments and Quasi-PDF
- Renormalization of quasi-PDF
- Conclusion

Conclusion

Using the operator product expansion, we proved the large-momentum factorization of the quasi-Parton Distribution Function. The matching formula between quasi-PDF and PDF is:

$$\tilde{q}\left(\tilde{x}, \frac{\mu^2}{(P^z)^2}, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \int_{-1}^1 \frac{dx}{|x|} C\left(\frac{\tilde{x}}{x}, \frac{\mu^2}{(x P^z)^2}\right) q\left(x, \frac{\mu}{\Lambda_{\rm QCD}}\right) + \mathcal{O}\left(\frac{\Lambda_{\rm QCD}^2}{(P^z)^2}\right)$$

or the coordinate space version:

$$\tilde{Q}\left(\zeta = P^{z} z, \frac{\mu^{2} \zeta^{2}}{(P^{z})^{2}}, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \int_{0}^{1} d\alpha \, \mathcal{C}\left(\alpha, \frac{\mu^{2} \zeta^{2}}{(P^{z})^{2}}\right) Q\left(\alpha \zeta, \frac{\mu}{\Lambda_{\rm QCD}}\right) + \mathcal{O}\left(\frac{\Lambda_{\rm QCD}^{2}}{(P^{z})^{2}}\right)$$

- OPE can be performed provided z^2 is small.
- At fixed P^z , in the $z^2 \to 0$ limit, only one operator, the local operator, will contributes: $\bar{\psi}(z)\gamma^z U(z,0)\psi(0) \approx C_0(\mu^2 z^2)\bar{\psi}(0)\gamma^z \psi(0)$. This can not be used to obtain PDF.
- However, in the $z^2 \rightarrow 0$ limit, if we keep $P^z z$ constant, then all the twist-2 operators will survive.

$$\tilde{Q}\left(\zeta = P^{z}z, \frac{\mu^{2}\zeta^{2}}{(P^{z})^{2}}, \frac{\mu}{\Lambda_{\rm QCD}}\right) = \sum_{n} C_{n}\left(\frac{\mu^{2}\zeta^{2}}{(P^{z})^{2}}\right) \frac{(i\,\zeta)^{n}}{n!} a_{n+1}\left(\frac{\mu}{\Lambda_{\rm QCD}}\right)$$
(63)

Thank You

Thank You!